



Nagoya City University Academic Repository

学位の種類	博士（医学）
報告番号	甲第1739号
学位記番号	第1236号
氏名	間所 佑太
授与年月日	令和2年3月25日
学位論文の題名	<p>Reduced Cholinergic Activity in the Hippocampus of Hippocampal Cholinergic Neurostimulating Peptide Precursor Protein Knockout Mice (HCNP 前駆体タンパクノックアウトマウスの海馬では、コリン作動性神経活動が減弱している)</p> <p>International Journal of Molecular Sciences, 28:101-107, 2019</p>
	<p>View metadata, citation and similar papers at CORE.ac.uk</p> <p>田口・ 幸平 佑太, 田川 誠</p> <p>provided by NCU Repository brought to you by CORE</p>

Abstract

The cholinergic efferent network from the medial septal nucleus to the hippocampus has an important role in learning and memory processes (Mesulam, et. al. *Neuroscience* 1983). This cholinergic projection can generate theta oscillations in the hippocampus to efficiently encode novel information (Dupret, et al. *Neuron* 2013). Hippocampal cholinergic neurostimulating peptide (HCNP) induces acetylcholine synthesis in medial septal nuclei (Ojika, et al. *Brain Res.* 1992). HCNP is processed from the N-terminal region of a 186 amino acid, 21 kD HCNP precursor protein called HCNP-pp (Ojika K, et al. *Prog. Neurobiol.* 2000) [also known as Raf kinase inhibitory protein (RKIP) and phosphatidylethanolamine-binding protein 1 (PEBP1)]. In this study, we generated HCNP-pp knockout (KO) mice and assessed their cholinergic septo-hippocampal projection, local field potentials in CA1, and behavioural phenotypes. No significant behavioural phenotype was observed in HCNP-pp KO mice. However, theta power in the CA1 of HCNP-pp KO mice was significantly reduced because of fewer cholineacetyltransferase-positive axons in the CA1 stratum oriens. These observations indicated disruption of cholinergic activity in the septo-hippocampal network. Our study demonstrates that HCNP may be a cholinergic regulator in the septo-hippocampal network.